

Latino Child Health: Need for Inclusion in the US National Discourse

ABSTRACT

The “rediscovery” of poverty, as echoed in concepts of social inequality, has contributed to the goal of eliminating racial/ethnic and social class disparities in the United States. This commentary focuses on what we know about the pressing health care needs and issues relevant to Latino children and families and how extant knowledge can be linked to priority policy recommendations to ensure the inclusion of Latino health issues in the national discourse.

A systematic review of the literature on Latino children and of expert opinion revealed 4 evidence-based themes focused on poverty: economic factors, family and community resources, health system factors, and pitfalls in Latino subgroup data collection. Consensus was found on 4 priority policy recommendations: (1) reduce poverty and increase access to health care coverage, (2) increase funding in targeted primary and preventive health care services, (3) provide funds needed to fully implement relevant health legislation, and (4) improve measurement and quality of data collection. If these recommendations are not instituted, the goals of Healthy People 2010 will not be achieved for the Latino population. (*Am J Public Health*. 2000;90:1827–1833)

Ruth E. Zambrana, PhD, and Laura A. Logie, BSEd

Past US federal initiatives designed to reduce racial and ethnic disparities have given limited attention to health inequities in the Latino population.^{1–3} An important question is the following: Given the historic presence of Latinos in the United States and the continuing increases in this population owing to population growth and immigration, why have Latinos failed to achieve inclusion in the national health policy agenda? There are several possible answers to this question.

First, Hispanics were not identified in the US census, on birth and death certificates, or in other national data systems until the past 2 decades.^{4–6} Second, presentation of health data indicating relatively good health outcomes in spite of low socioeconomic status for Hispanics in the aggregate obscured important differences by subgroup, generation, birthplace, and time in the United States. The application of the term *epidemiological paradox* to findings of good health outcomes explained the Mexican-origin outcomes^{7–12} but masked the heterogeneity of the Latino population. It also drew attention away from Latinos’ limited access to care and disproportionate burden of chronic diseases such as adult diabetes and childhood asthma.

Third, Latino scholars and federal and local officials have been relatively absent at the national, state, and local levels. National Latino organizations focused on Latino health did not emerge until the late 1990s, and this resulted in a lack of advocacy at the national level. Finally, a limited empirical base was available to support claims for the disproportionate burdens of death and disability that constituted principal criteria for inclusion in the national health discourse.

Social and Economic Equality?

Healthy People 2010,² the national disease prevention and health promotion goals for the year 2010, has the overarching aim of elim-

inating disparities. For the first time, objectives have been set and baseline and monitoring data are presented for Latino subgroups.^{13,14} The goal of eliminating racial/ethnic disparities requires a considerable scholarly investment if the reasons underlying disparities are to be understood; there has not been adequate research on inclusion and measurement of medical and nonmedical factors associated with Latino child and family health outcomes. Herein resides the concern that goals may not be achieved unless changes are implemented within a broader framework of social and economic equity and community resource development to improve Latino health.^{15–19}

This commentary focuses on what we know about the pressing health care needs and issues relevant to Latino child and family health and how extant knowledge can be linked to priority policy recommendations to ensure the inclusion of Latino health in the national discourse. Although we focus on children, we do so in the context of their families.

Why Latino Children? Why Now?

Growth rates among the Latino population not only exceed those of other minority groups but also exceed the growth rate of the US population as a whole.^{20,21} If projected growth trends continue, the Latino population will be the largest minority group in the United States, reaching a projected 96.5 million people (24.5% of the total US population) within the next 5 years.^{22,23} Relative to Whites and African Americans, Latinos are a young popula-

The authors are with the Department of Women’s Studies, University of Maryland College Park.

Requests for reprints should be sent to Ruth E. Zambrana, PhD, Department of Women’s Studies, University of Maryland, 2101 Woods Hall, College Park, MD 20742 (e-mail: rz23@umail.umd.edu).

This commentary was accepted August 15, 2000.

TABLE 1—Socioeconomic and Health Indicators for US Latino Children

	Latino, %	Black, %	Non-Hispanic White, %
Socioeconomic data			
Children in US population	15	15	65
Children in poverty	36	37	11
Families with children younger than 18 years living at home	64	36	76
Children in single-parent homes	31	55	21
Health indicators			
Access/use of services			
Children without usual source of care	17.2	12.6	6
Children without health insurance	27.7	17.6	12.2
Children with Medicaid coverage	32	37	14
Children with private health insurance	43	47	79
Families "unable to afford health care"	69.1	60.4	58.5
Children "in excellent health"	42.9	48.1	55.3
Children aged 1–4 years with no physician visit	11.7	11.1	8.4
Children aged 5–9 years with no physician visit	24.4	18.3	17.9
Youth aged 10–14 years with no physician visit	34.2	28.9	24.8
Adolescents aged 15–19 years with no physician visit	37	26.7	25.7
Risk behaviors			
12th graders reporting heavy drinking	14	8	27
12th graders reporting illicit drug use	24	20	27
10th graders reporting smoking cigarettes daily	9.4	5.8	20.3
Female youth reporting use of condom at most recent intercourse	40	58.9	49.2
Male youth using condom inconsistently or not at all	71	53	54
Teen births	17.4	22.8	11.3
High school girls attempting suicide	14.9	9	10.3
Preventive health behaviors			
Children not fully immunized	24	23	20
Children with active asthma	11 ^a	6	3
Children aged 2–4 years with dental caries in primary teeth	32	22	13
Children aged 5–17 years with dental caries in permanent teeth	49	39	45
Boys aged 6–11 years who are overweight	18.8	14.7	13.2
Girls aged 6–11 years who are overweight	15.8	17.9	11.9

^aPuerto Ricans.

tion. Approximately 35% of Latinos (vs 24% of Whites and 30% of African Americans) are younger than 18 years.²²

The number of Latino children has already surpassed that of any other minority group, growing from 9% of the child population in 1980 to 15% in 1999. In 1998, the number of Latino children was estimated to be 10.5 million.²³ Although only 13% of Latino children were foreign born in 1996, 31% report having difficulty speaking English. By the year 2020, an estimated 1 in 5 children living in the United States will be Latino.²⁰

Latino children account for a large and growing segment of the nation's student population (13.5%). States with high Latino elementary and secondary school enrollments include New Mexico (46.6%), California (38.7%), Texas (36.7%), Arizona (30%), Colorado (18.4%), New York (17.4%), Florida (15.3%), New Jersey (13.5%), Illinois (12.2%), and Massachusetts (9.3%).²² The aforementioned demographic changes provide a compelling rationale for federal and state policymakers to include the health care needs of Latino children in their health policy agendas to achieve the Healthy People 2010 objective of eliminating ethnic disparities.

Table 1 displays selected socioeconomic and health indicators for Latino children in

comparison with African American and non-Hispanic White children. Latino children are disproportionately represented among the poor in the United States. In 1998, 34.4% of Latino children lived in poverty, although there were significant differences among subgroups. Puerto Rican children (43.5%) are most likely to live in poverty, followed by Mexican American (35.4%), Central and South American (26.6%), and Cuban (16.4%) children.²²

Latino children are more likely to be uninsured than African American or non-Hispanic White children.^{19,24–29} Mexican Americans are the Latino group most likely to be uninsured.³⁰ High rates of noncoverage are associated with the following 2 factors: parents working in economic sectors that lack employment-linked health benefits^{19,24,28,31,32} and multiple and persistent barriers experienced by Latinos in accessing health care.^{27,29,33–39} More than one third (36%) of Medicaid-eligible Latino children are not enrolled, and only about 32% of eligible Latino children receive Medicaid benefits.^{35,37,40,41} Among Latino children, lack of health insurance is strongly associated with greater use of emergency rooms as a source of primary care,^{33,36,42} less likelihood of having a usual source of care,^{29,34,37,43} and less contact with a physician in

the previous year.^{44,45} The older the Latino child, the more likely he or she is to have no physician contact relative to his or her African American and non-Hispanic White peers.

Available evidence shows that in comparison with African American and non-Hispanic White children, Latino children generally (1) are less likely to be immunized,^{43,46–51} (2) have higher rates of tuberculosis,^{52,53} (3) have higher rates of obesity and sedentary activity,^{54–59} (4) have more dental caries,^{43,57} (5) are more likely to experience intentional and unintentional injuries,^{60–62} and (6) are more likely to reside in hazardous environments.^{21,43,63–65} Latino adolescents are more likely to use drugs, alcohol, and tobacco^{21,43,66}; less likely to use contraceptives^{67,68}; more likely to be injured^{21,43,66}; and more likely to attempt suicide than African American and non-Hispanic White adolescents.^{69–71}

Evidence-Based Themes: What Do We Know About Latino Child and Family Health?

Four major themes emerged from the body of evidence reviewed^{72–74}: (1) the role of socioeconomic status and its strong associa-

tion with health insurance coverage and adverse health outcomes^{15,19,24,25,27–29,31,35,40,65,75,76}; (2) the importance of family and community resources in improving Latino child health^{76–97}; (3) health system factors as key predictors of use of services, compliance behaviors, and health outcomes^{33,34,36,47,52,98–104}; and (4) the recognition that Latinos are not a homogeneous group.^{105–113}

Economic Factors

Economic factors as predictors of health access and outcomes were notably more likely to be included in post-1991 investigations of Latino health.^{18,34,35,40,41,48,75,76} In comparison with African Americans and non-Hispanic Whites, Latinos are more likely to report that they are unable to afford health insurance, more likely to work in employment sectors that do not have employment-linked health benefits, and less likely to access public benefits such as Medicaid.^{19,24,27,31,35,40} Although language is a barrier to access, ethnic disparities have been shown to persist even after limited English-language proficiency has been taken into account.^{21,42,101,114}

Several recent reviews of the literature on access among Latinos substantiate the links between socioeconomic status and health.^{3,15,24–26,28,29,46,76,115} Low socioeconomic status is a significant determinant of health status and outcomes for Latino children and accounts for a substantial proportion of ethnic-specific group disparities in morbidity and mortality patterns.^{24–26,28,32,76,116,117}

Family and Community Resources

Familial behaviors and community resources have been identified as key factors in promoting children's physical and mental well-being.^{76–97} Parental factors, particularly education and literacy levels, play a direct role in Latino children's health access and outcomes.^{39,47,52,61,96,98} Families, often not aware of services or fearful of providers, place Latino children at risk for less use of preventive screening services, more lost school days, and potential health problems in adulthood.^{4,29,91,100} Providers' involvement with parents may significantly increase the use of preventive and primary care services.^{36,38,92}

Resources available in a community are linked with availability of health care services and are powerful determinants of use of preventive and primary care services and of health outcomes.^{15,34,35,41,47,77,78,118} Health care services, often overburdened or entirely lacking in low-income Latino communities, are not able to adequately respond to the needs of Latino children.^{15,76} Therefore, working in partnership with the community and its families to develop ethnic-specific clinical screening services, parent education, and outreach programs can help improve Latino children's

health.^{77,78,80,81,83,85,88,89,91,92,94,96,98} School health clinics, community prevention efforts, disease-centered outreach programs, and pediatric clinics are some of the innovative interventions that have proven effective.^{36,41,77–80,82,84,86,87,95,97}

Health System Factors

Institutional systems and barriers are consistently identified as powerful predictors of parental health behaviors, use of services, and child health outcomes.^{33,34,36,47,52,98–104} Institutional factors include provider attitudes, availability of translators, waiting times, location, and other nonfinancial barriers.^{27,36} Access to the health care system does not ensure that appropriate services will be provided or that a child's health will improve. Inequities in use of preventive and primary care services have been documented with respect to vision screening, prescription medications and equipment for the management of asthma, and access to and use of mental health services.^{31,33,38,98,99,116,117} Changes in health system factors can best be achieved by implementing and monitoring the US Department of Health and Human Services cultural competency guidelines as well as civil rights guidance for limited English proficiency (Executive Order 13166, issued on August 11, 2000), and by increasing the numbers of bilingual and bicultural providers.^{13,14,118}

Pitfalls in Latino Subgroup Data Collection

Authors have consistently observed that health outcomes can be explained by subgroup differences.^{105–113} Although significant progress has been made in the identification and collection of data on Latino health, several methodologic limitations hinder researchers' ability to fully capture disease and disability trends among Latino subgroups.^{6,13,14,119} These limitations include the following: aggregation of all Hispanics into one group for analysis, without precise measurement of socioeconomic status; limited reporting of age and sex breakdowns, especially in studies of Latino children; and exclusion of Spanish-speaking populations, predominantly immigrants, because most national survey instruments are published in English.^{4,5,13,14,118–121} Current federal and state data systems are also limited in terms of Latino child health indicators.⁶⁴

A systematic review of approximately 200 studies on Latino children revealed several methodologic limitations: more than 60% of the studies did not report a method of ethnic identification; most reported data for Hispanics as an overall group; and almost half did not report an indicator for socioeconomic status. Also, because more than 50% of the studies involved secondary data, there was often an absence of relevant med-

ical and nonmedical factors to explain the findings.

Most research has focused on Mexican-origin children, with an emphasis on nutrition, obesity, and physical fitness. Such studies often cite data derived from the Hispanic Health and Nutrition Examination Survey^{39,50,113,122–126} or the third National Health and Nutrition Examination Survey, which included only Mexican Americans.^{55,56,127,128} A recent review of Latino child health revealed that major research gaps include lack of subgroup data on parental and child health behaviors, chronic childhood illnesses, intentional and unintentional injuries, environmental health, migrant and US–Mexico border resident health, and mental health.¹⁰¹

Measurement of a broader array of factors, such as community context, parental and child health behaviors, and other lifestyle factors, is important for understanding the links between poverty and adverse health outcomes.^{16,17,39,76–80,96,101,117} Broadening the scientific paradigm for understanding Latino child health involves examining the links among family, community, and cultural protective factors and institutional factors.^{49,56,77,80,84,101,117} Although data limitations and research gaps have impeded the strategic process of including the needs of Latino children in the national health discourse, sufficient evidence now exists to inform priority policy recommendations.^{29,102,129–131}

Policy Recommendations

A panel of 7 experts on Latino child health convened for a 2-day meeting in July 1999 to review what is known about Latino family and child health and to link the evidence to priority policy recommendations. The experts forged a consensus on 4 priority policy areas for promoting long-term, favorable Latino child health outcomes (see Table 2). These policy issues were as follows.

1. Decreasing rates of poverty among Latino families and children call for legislation and policies that promote economic equity by guaranteeing a minimum living wage, providing employment-linked health benefits,^{19,28} and increasing access to adult education and literacy programs.

2. The most pressing policy area for Latino families and children is lack of access to health care.^{15,24,25,27,29,32–37,39} Expanding access to health care services for Latino children and their families requires raising eligibility cutoffs to 250% of the poverty level for Medicaid, the State Child Health Insurance Program, and prenatal care programs for poor and working-class Latino families, especially those without employee-linked health benefits. Although many Latino families are eligible for such services, enrollment is low.^{28,41,132,133} For example, Texas has extremely low Medicaid

TABLE 2—Evidence-Linked Findings and Priority Policy Recommendations: July 1999

Finding	Recommendations
Socioeconomic status is significantly linked to lack of health insurance	<p>Decrease rates of poverty</p> <p>Guarantee minimum living wage</p> <p>Provide employment-linked health benefits</p> <p>Provide access to adult education and literacy programs</p> <p>Increase access to health care</p> <p>Raise eligibility to 250% of poverty level for federal and state programs</p> <p>Increase funding for outreach to promote awareness of programs & enrollment</p> <p>Monitor compliance with public charge guidance</p>
Families & community resources are key factors for improving child health	<p>Increase funding for primary and preventive care services targeting younger children</p> <p>Address high-priority areas of health</p> <p>Increase funding for school and mental health services</p>
Health system factors are important predictors of health behaviors, use of services, and health outcomes	<p>Provide sufficient allocations of funds to fully implement:</p> <p>Disadvantaged Minority Health Improvement Act</p> <p>Health Professions Partnership Act</p> <p>Cultural competency guidelines</p> <p>Civil rights guidance for limited English proficiency</p>
Latinos are not a homogeneous group	<p>Improve measurement and quality of data collection</p> <p>Strengthen quality of data in state vital statistics registries</p> <p>Implement standard administration of surveys in Spanish</p> <p>Report national and state data by at least 3 homogeneous subgroups</p>

eligibility cutoffs, and a large number of Latino children remain uninsured.¹³³ To increase enrollment, additional funding is needed for informational outreach on eligibility via Spanish-language media and at sites such as public health clinics, schools, and churches. In addition, compliance with US public charge regulations is recommended to ensure that eligible legally-resident families enrolling in the State Child Health Insurance Program, Medicaid, or other noncash benefit programs will not jeopardize their immigrant status.^{76,83,91,94–96,134}

3. Directly associated with documented health system inequities is the recommendation to increase federal and state funding for research and targeted preventive and primary care interventions that address identified health priority concerns of Latino children and families.^{31,33,38,98,99,116,117} Family-focused primary and preventive health care services located in the community (with schools as important “windows of clinical opportunity”) can effectively decrease health risks for children and families.^{80–87,91–97} Ethnic-specific preventive and primary care services must be accompanied by sufficient authorization of funds to fully implement the 1990 Disadvantaged Minority Health Improvement Act, the Health Professions Partnership Act of 1998, and civil rights guidance for limited English proficiency.^{3,6,13,14,118,121} Focused investments for the education, training, and hiring of bilingual and bicultural providers, particularly for the historically underrepresented Latinos, and enforcement of compliance with the Department

of Health and Human Services cultural competency guidelines can significantly contribute to the aim of eliminating ethnic disparities.¹¹⁸

4. Although significant progress has been made in data collection, problems persist in the measurement and quality of data collected. To ensure the development of more comprehensive baseline indicators and the monitoring and surveillance of health indicators for Latinos, public health data systems must be improved through collection of minimal core information on racial/ethnic subgroups, place of birth, education, and English-language proficiency levels; administration of surveys in Spanish to ensure representation and inclusion of Latinos who have difficulty with English; and reporting of national and state data for at least 3 major Latino subgroups.^{4,5,6,12–14,26,40,64,119,135,136} These data can yield community-specific baseline indicators. In 1998, the National Center for Health Statistics began the administration of Spanish-language instruments for the National Health Interview Survey; bilingual instruments should routinely be used in public health data collection procedures.⁴ Progress in collecting baseline data is particularly important, as seen in the fact that 21% of the goals of Healthy People 2000 were not assessable.¹¹⁵

Conclusions

These policy recommendations derived from a literature review and from expert opin-

ions show that health equity cannot be achieved without simultaneously improving economic equity to offset the effects of poverty on health outcomes for Latino children.^{16,18} Current state programs such as the Children’s Health Insurance Program and Medicaid access to prenatal care programs^{132,137} can most effectively address access to care for low-income and working poor Latino families and children. The literature also reveals that access to clinical preventive screening and primary care services is necessary to reduce inequities in Latino child health.

These policy recommendations are neither new nor innovative, but they confirm those made in the past^{1,120,121,138,139} and are consistent with recent federal initiatives.^{13,14,118} In answering the question of why Latinos have failed to achieve inclusion in the health policy agenda, we suggest that past lack of political representation, combined with methodologic limitations, has significantly contributed to the exclusion of Latino health from the national discourse.

Recent reports such as *Hispanic Agenda for Action*¹²¹ and *Healthy People 2000 Progress Review*,¹¹⁵ along with the 1999 health workshop of the Hispanic Congressional Caucus Institute, reinforce policy priorities similar to those described here. Our data provide additional evidence of a compelling scientific consensus on the public health needs and priority policy recommendations necessary to ensure social and economic equity and reduce health disparities for low-income Latino children and their families. □

Contributors

R.E. Zambrana planned, designed, and conceptualized the study and wrote the paper. L.A. Logie contributed to the conceptualization and writing of the paper.

Acknowledgments

The Robert Wood Johnson Foundation (grant 37533) provided financial support for this project.

We are grateful for the thoughtful contributions of Olivia Carter-Pokras, PhD, Division of Policy and Data, Office of Minority Health; Glen Flores, MD, Department of Pediatrics and Public Health, Boston University School of Medicine; and the anonymous reviewers. We would like to thank Darrell J. Gaskin, PhD, Georgetown University, for his help in the data analysis.

We also deeply appreciate the contributions of our panel of experts in Latino Child Health, including Glenn Flores, MD, Boston University Schools of Medicine and Public Health; Olivia Carter-Pokras, PhD, Division of Policy and Data, Office of Minority Health, US Department of Health and Human Services; Fernando Mendoza, MD, MPH, Department of Pediatrics, Stanford University School of Medicine; Robert O. Valdez, PhD, Department of Health Policy and Management, University of California, Los An-

geles; Virginia Bishop-Townsend, MD, MPH, Children's Memorial Hospital, Chicago, Ill; Hector Balcazar, PhD, Center of Behavioral and Social Sciences, School of Public Health, University of North Texas; and Hilda Ochoa Bogue, MS, CHES, Migrant Health Services, Community Health of South Dade, Florida.

Finally, we acknowledge the health staff of the National Council of La Raza, especially Norma Y. Lopez, vice president for health promotion and health policy, for their commitment to improving Latino health.

References

1. Heckler M. *Report of the Secretary's Task Force on African American and Minority Health. Vol I: Executive Summary*. Washington, DC: US Department of Health and Human Services; 1985.
2. *Healthy People 2010*. Washington, DC: US Department of Health and Human Services; 2000.
3. *Eliminating Racial and Ethnic Disparities in Health*. Washington, DC: US Department of Health and Human Services, Office of Minority Health; 1999.
4. Hajat A, Lucas J, Kington R. Health outcomes among Latino subgroups: United States, 1992–95. *Adv Data Vital Health Stat*. February 25, 2000; No. 310. DHHS publication PHS 2000-1250.
5. Zambrana RE, Carter-Pokras O. Health data issues for Latinos: implications for public health research. *J Health Care Poor Underserved*. In press, February 2001.
6. Public Health Service Task Force on Minority Health Data. *Improving Minority Health Statistics*. Washington, DC: US Public Health Service; 1992.
7. Abraido-Lanza AF, Dohrenwend B, Ng-Mak D, Turner J. The Latino mortality paradox: a test of the "salmon bias" and healthy migrant hypotheses. *Am J Public Health*. 1999;89:1543–1548.
8. Fuentes-Afflick E, Hessel NA, Perez-Stable EJ. Testing the paradox of low birth weight in Latinos. *Arch Pediatr Adolesc Med*. 1999;153:147–153.
9. Markides KS, Coreil J. The health of Latinos in the southwestern United States: an epidemiologic paradox. *Public Health Rep*. 1986;101:253–265.
10. Scribner R. Infant mortality among Latinos: the epidemiological paradox. *JAMA*. 1991;265:2065–2066.
11. Scribner R. Paradox as paradigm—the health outcomes of Mexican Americans. *Am J Public Health*. 1996;86:303–305.
12. Balcazar H, Aoyama C, Cai X. Interpretative views on Hispanics' perinatal problems of low birth weight and prenatal care. *Public Health Rep*. 1991;106:420–426.
13. *Setting a Research Agenda: Challenges for the Minority Health Statistics Grants Program*. Hyattsville, Md: National Center for Health Statistics; 1991. Available at: www.cdc.gov/nchs/about/grants/minpop/Hispanic/Hispanic.htm. Accessed October 12, 2000.
14. Ad Hoc Working Group on Racial and Ethnic Data. *Meeting Current and Future Demands for Racial and Ethnic Data for Health and Human Services*. Rockville, Md: US Department of Health and Human Services; 1996.
15. Smith MW, Kreutzer RA, Goldman L, Casey-Paál A, Kizer KW. How economic demand influences access to medical care for rural Hispanic children. *Med Care*. 1996;34:1135–1148.
16. Pappas G, Quenn S, Hadden W, Fisher G. The increasing disparity in mortality between socioeconomic groups in the United States, 1960 and 1986. *N Engl J Med*. 1993;329:103–109.
17. Rumbaut R, Weeks JR. Unraveling a public health enigma: why do immigrants experience superior perinatal health outcomes? *Res Soc Health Care*. 1996;13B:337–391.
18. Adler NE, Boyce T, Chesney MA, Folkman S, Syme L. Socioeconomic inequalities in health: no easy solution. *JAMA*. 1987;269:3140–3145.
19. Santos R, Seitz P. Benefit coverage for Latino and Latina workers. In: Perez SM, ed. *Moving Up the Ladder: Latino Workers and the Nation's Future Prosperity*. Washington, DC: National Council of La Raza; 2000:162–185.
20. *Changing America: Indicators of Social and Economic Well-Being by Race and Hispanic Origin*. Washington, DC: Council of Economic Advisors for the President's Initiative on Race; 1998.
21. *Trends in the Well-Being of America's Children and Youth 1998*. Washington, DC: US Department of Health and Human Services, Office of the Assistant Secretary for Planning and Evaluation; 1998.
22. Ramirez RR. *The Hispanic Population in the United States: Population Characteristics*. Washington, DC: US Bureau of the Census; 2000. Current Population Report 20-527.
23. National Council of La Raza. Demographic snapshot of the Hispanic population. *Agenda Newsletter*. 1998;14:1.
24. Zuvekas SH, Weinick RM. *Changes in Access to Care, 1977–1996: The Role of Health Insurance*. Washington, DC: US Department of Health and Human Services; 1999. DHHS publication 99-R054.
25. Simpson G, Bloom B, Cohen RA, Parsons PE. Access to health care, I: children. *Vital Health Stat 10*. 1997;No. 196:1–46.
26. Coiro MJ, Zill N, Bloom B. Health of our nation's children. *Vital Health Stat 10*. 1994; No. 191:1–61.
27. Cornelius LJ. Barriers to medical care for white, black and Hispanic American children. *J Natl Med Assoc*. 1993;85:281–288.
28. *Working Without Benefits: The Health Insurance Crisis Confronting Hispanic Americans*. New York, NY: Commonwealth Fund; 2000.
29. *No Health Insurance? It's Enough to Make You Sick*. Philadelphia, Pa: American College of Physicians–American Society of Internal Medicine; 2000.
30. Holl JL, Szilagyi PG, Rodewald LE, Byrd RS, Weitzman ML. Profile of uninsured children in the United States. *Arch Pediatr Adolesc Med*. 1995;149:398–406.
31. Braveman P, Olivia G, Miller MG, et al. Adverse outcomes and lack of health insurance among newborns in an eight-county area of California. *N Engl J Med*. 1989;321:508–513.
32. Andersen R, Lewis SZ, Giachello AL, Aday LA, Chiu G. Access to medical care among the Hispanic population of the southwestern United States. *J Health Soc Behav*. 1981;22:78–89.
33. Davidson AB, Klein DB, Settignano GA, Alano AJ. Access to care among children visiting the emergency room with acute exacerbations of asthma. *Ann Allergy*. 1994;72:469–473.
34. Lieu TA, Newacheck PW, McManus MA. Race, ethnicity, and access to ambulatory care among US adolescents. *Am J Public Health*. 1993;83:960–965.
35. Halfon N, Wood D, Valdez R, Burciaga R, Pereyra M, Duan N. Medicaid enrollment and health services access by Latino children in inner-city Los Angeles. *JAMA*. 1997;277:636–641.
36. Flores G, Vega L. Barriers to health care access for Latino children: a review. *Fam Med*. 1998;30:196–205.
37. Bartman BA, Moy E, D'Angelo LJ. Access to ambulatory care for adolescents: the role of a usual source of care. *J Health Care Poor Underserved*. 1997;8:214–226.
38. Lara M, Allen F, Lange L. Physician perceptions of barriers to care for inner-city Latino children with asthma. *J Health Care Poor Underserved*. 1999;10:27–44.
39. Estrada AL, Trevino FM, Ray LA. Health care utilization barriers among Mexican Americans: evidence from HHANES 1982–84. *Am J Public Health*. 1990;80(suppl):27–31.
40. Avruch S, Machlin S, Bonin P, Ullman F. The demographic characteristics of Medicaid-eligible uninsured children. *Am J Public Health*. 1998;88:445–447.
41. Short PF, Lefkowitz DC. Encouraging preventive services for low-income children. The effect of expanding Medicaid. *Med Care*. 1992;30:766–780.
42. Zambrana RE, Ell K, Dorning C, Wachsmann L, Hodge D. The relationship between psychosocial status of immigrant Latino mothers and use of pediatric emergency services. *Health Soc Work*. 1994;19:93–102.
43. *America's Children: Key National Indicators of Well-Being, 2000*. Washington, DC: Federal Interagency Forum on Child and Family Statistics; 2000.
44. Butler JA, Singer JD, Palfrey JS, Walker DK. Health insurance coverage and physician use among children with disabilities: findings from probability samples in five metropolitan areas. *Pediatrics*. 1987;79:89–98.
45. *Child Health USA 1999*. Washington, DC: Health Resources and Services Administration; 1999.
46. Centers for Disease Control and Prevention. Vaccination coverage by race/ethnicity and poverty level among children aged 19–35 months—United States, 1996. *MMWR Morb Mortal Wkly Rep*. 1996;46:963–969.
47. Wood D, Donald-Sherbourne C, Halfon N, et al. Factors related to immunization status among inner-city Latino and Mexican-American preschoolers. *Pediatrics*. 1995;96:295–301.
48. Mayer ML, Clark SJ, Konrad TR, Freeman VA, Slifkin RT. The role of state policies and programs in buffering the effects of poverty on children's immunization receipt. *Am J Public Health*. 1999;89:164–170.
49. Guendelman S, English P, Chavez G. The effects of maternal health behaviors and other risk factors on immunization status among Mexican-American infants. *Pediatrics*. 1995;95:823–828.
50. Gergen PJ, Ezzati T, Russell H. DTP immunization status and tetanus antitoxin titers of Mexican American children ages six months

- through eleven years. *Am J Public Health*. 1988; 78:1446–1450.
51. Moore P, Fenlon N, Hepworth JT. Indicators of differences in immunizations of Mexican American and white non-Hispanic infants in a Medicaid managed care system. *Public Health Nurs*. 1996;13:21–30.
52. Watchi R, Kahlstrom E, Vachon LA, Barnes PF. Pediatric tuberculosis: clinical presentation and contact investigation at an urban medical center. *Respir*. 1998;65:192–194.
53. Ussery XT, Valway SE, McKenna M. Epidemiology of tuberculosis among children in the United States: 1985 to 1994. *Pediatr Infect Dis J*. 1996;15:697–704.
54. Alexander MA, Sherman JB, Clark L. Obesity in Mexican-American preschool children—a population group at risk. *Public Health Nurs*. 1991;8:53–58.
55. Centers for Disease Control and Prevention. Update: prevalence of overweight among children, adolescents, and adults—United States, 1988–1994. *MMWR Morb Mortal Wkly Rep*. 1997;46:198–202.
56. Andersen RE, Crespo CJ, Bartlett SJ, et al. Relationship of physical activity and television watching with body weight and level of fatness among children: results from the third National Health and Nutrition Examination Survey. *JAMA*. 1998;279:938–942.
57. *Health, United States, 2000 With Adolescent Health Chartbook*. Hyattsville, Md: National Center for Health Statistics; 2000.
58. Finkenberger MB, DiNucci JM. Age, ethnic and gender differences in physical fitness of middle-school children in east Texas. *Percept Mot Skills*. 1995;80:387–393.
59. Guinn B, Semper T, Crofts A. Anthropometric comparisons among low-income Mexican-American children aged 10 through 14. *J Am Diet Assoc*. 1986;86:57–60.
60. Baker SP, Braver ER, Chen LH, Pantula J, Massie D. Motor vehicle deaths among Latino and African American children and teenagers. *Arch Pediatr Adolesc Med*. 1998;152:1209–1212.
61. Mull DS, Agran PF, Winn DG, Anderson CL. Household poisoning exposure among children of Mexican-American born mothers: an ethnographic study. *West J Med*. 1999;171:16–19.
62. Matteucci RM, Holbrook TL, Hoyt DB, Molgaard C. Trauma among Hispanic children: a population-based study in a regionalized system of trauma care. *Am J Public Health*. 1995;85:1005–1008.
63. Mott L. The disproportionate impact of environmental health threats on children of color. *Environ Health Perspect*. 1995;103(suppl 6):33–35.
64. Metzger R, Delgado JL, Herrell R. Environmental health of Latino children. *Environ Health Perspect*. 1995;103(suppl 6):25–32.
65. Wernette DR, Nieves LA. Breathing polluted air: minorities are disproportionately exposed. *EPA J*. 1992;18:16–17.
66. Johnston LD, O'Malley PM, Bachman KG. *National Survey Results on Drug Use From the Monitoring the Future Study, 1975–1997*. Rockville, Md: National Institute on Drug Abuse; 1998. NIH publication 98-4345.
67. Marcell A. Understanding ethnicity, identity formation and risk behavior among adolescents of Mexican descent. *J Sch Health*. 1994;64:8–15.
68. Mayden B, Castro W, Annetto M. *First Talk: A Teen Pregnancy Prevention Dialogue Among Latinos*. Washington, DC: Child Welfare League of America Press; 1999.
69. Hovey JD, King CA. Acculturative stress, depression, and suicidal ideation among immigrant and second-generation Latino adolescents. *J Am Acad Child Adolesc Psychiatry*. 1996;35:1183–1192.
70. Queralto M. Risk factors associated with completed suicide in Latino adolescents. *Adolescence*. 1993;28:831–850.
71. Roberts RE, Chen YR, Roberts CR. Ethnocultural differences in prevalence of adolescent suicidal behaviors. *Suicide Life Threat Behav*. 1997;27:208–217.
72. Briss PA, Zaza S, Pappaioanu M, et al. Developing an evidenced-based guide to community preventive services—methods. *Am J Prev Med*. 2000;18(suppl 1):35–43.
73. Cooper H, Hedges LV, eds. *The Handbook of Research Synthesis*. New York, NY: Russell Sage Foundation Press; 1994.
74. Cook TD, Cooper H, Cordray DS, et al. *Meta-Analysis for Explanation—A Casebook*. New York, NY: Russell Sage Foundation Press; 1992.
75. Valdez RB, Morgenstern H, Brown R, Wyn R, Wang C, Cumberland W. Insuring Latinos against the cost of illness. *JAMA*. 1993;269:889–894.
76. Bagley SP, Angel R, Dilworth-Anderson P, Liu W, Schinke S. Panel V: adaptive health behaviors among ethnic minorities. *Health Psychol*. 1995;14:632–640.
77. Brindis C. Adolescent pregnancy prevention for Hispanic youth: the role of schools, families, and communities. *J Sch Health*. 1992;62:345–351.
78. Waterman SH, Hill LL, Robyn B, et al. A model immunization demonstration for preschoolers in an inner-city barrio, San Diego, California, 1992–1994. *Am J Prev Med*. 1996;12(suppl 4):8–13.
79. Setzer JR, Smith DP. Comprehensive school based services for pregnant and parenting adolescents in West Dallas, Texas. *J Sch Health*. 1992;62:97–102.
80. Flores G, Abreu M, Philipp BL, et al. Addressing persistent pockets of need for childhood immunization: use of an urban drop-in vaccination clinic by high-risk children. *Ambulatory Child Health*. 1999;5:3–13.
81. Kasper MJ, Robbins L, Root L, Peterson MGE, Allegrante JP. A musculoskeletal outreach screening treatment and education program for urban minority children. *Arthritis Care Res*. 1993;6:126–133.
82. McGurk SR, Cárdenas J, Adelman HS. Utilization of a school-based clinic for identification and treatment of adolescent sexual abuse. *J Adolesc Health*. 1993;14:196–201.
83. Mendoza FS, Fuentes-Afflick E. Latino children's health and the family-community health promotion model. *West J Med*. 1999;170:85–92.
84. Centers for Disease Control and Prevention. State-specific pregnancy and birth rates among teenagers—United States, 1991–1992. *MMWR Morb Mortal Wkly Rep*. 1995;44:677–683.
85. Agran PF, Winn DG, Anderson CL, Del Valle CP. Pediatric injury hospitalization in Hispanic children and non-Hispanic white children in Southern California. *Arch Pediatr Med*. 1996;150:400–406.
86. Birrer RB, Fleisher JM, Cortese L, et al. An urban primary school health program. *NY State J Med*. 1991;91:339–341.
87. Denison AV, Pierce JR. Enrollment in English-as-a-second-language class as a predictor of tuberculosis infection in schoolchildren. *Public Health Rep*. 1996;111:428–430.
88. Fitzgibbon ML, Stolley MR, Avellone MB, Sugerman S, Chavez N. Involving parents in cancer risk reduction: a program for Hispanic American families. *Health Psychol*. 1996;15:413–422.
89. Frauenglass S, Routh DK, Pantin HM, Mason CA. Family support decreases influence of deviant peers on Hispanic adolescents' substance use. *J Clin Child Psychol*. 1997;26:15–23.
90. Higgins PG, Burton M. New Mexico women with no prenatal care: reasons, outcomes, and nursing implications. *Health Care Women Int*. 1996;17:255–269.
91. Hubbell F, Waitzkin H, Mishra SI, Dombrink J. Evaluating health-care needs of the poor: a community-oriented approach. *Am J Med*. 1989;87:127–131.
92. Lewis MA, Lewis CB, Leake B, Monahan G, Rachelefsky G. Organizing the community to target poor Latino children with asthma. *J Asthma*. 1996;33:289–297.
93. McFarlane J. De Madres a Madres: an access model for primary care. *Am J Public Health*. 1996;86:879–880.
94. Nader PR, Sallis JF, Patterson TL, et al. A family approach to cardiovascular risk reduction: results from the San Diego Family Health Project. *Health Educ Q*. 1989;16:229–244.
95. Perez MA, Pinzon HL, Garza RD. Latino families: partners for success in school settings. *J Sch Health*. 1997;67:182–184.
96. Vega WA, Sallis JF, Patterson T, Rupp J, Atkins C, Nader PR. Assessing knowledge of cardiovascular health-related diet and exercise behaviors in Anglo- and Mexican-Americans. *Prev Med*. 1987;16:696–709.
97. Walter HJ, Vaughan RD, Armstrong B, Krakoff RY, Tiezzi L, McCarthy JF. School-based health care for urban minority junior high school students. *Arch Pediatr Adolesc Med*. 1995;149:1221–1225.
98. Wasserman RC, Croft CA, Brotherton SE. Preschool vision screening in pediatric practice: a study from the Pediatric Research in Office Settings (PROS) Network. *Pediatrics*. 1992;89:834–838.
99. Hahn BA. Children's health: racial and ethnic differences in the use of prescription medications. *Pediatrics*. 1995;95:727–732.
100. Chavez GF, Ellis AA. Pediatric hospital admissions for measles. Lessons from the 1990 epidemic. *West J Med*. 1996;165:20–25.
101. Flores G, Zambrana RE. The early years: the health of children and youth. In: Aguirre-Molina M, Molina C, Zambrana RE, eds. *Health Issues in the Latino Community*. San Francisco, Calif: Jossey-Bass. In press, Spring 2001.
102. Guendelman S. Developing responsiveness to the health needs of Hispanic children and families. *Soc Work Health Care*. 1983;8:713–730.
103. Amaya M, Ackall G, Pingitore N, Quiroga M, Terrazas-Ponce B. Childhood lead poisoning on the US-Mexico border: a case study in environ-

- mental health. *Public Health Nurs.* 1997;14: 353–360.
104. Waldman HB. Evolving “faces” of the next generation of pediatric patients. *J Dent Child.* January/February 1997;39–42.
105. Centers for Disease Control and Prevention. Childbearing patterns among Puerto Rican Hispanics in New York City and Puerto Rico. *MMWR Morb Mortal Wkly Rep.* 1987;36:39–41.
106. Bass JL, Mehta KA, Eppes B. Parasitology screening of Latin American children in a primary care clinic. *Pediatrics.* 1992;89:279–283.
107. Becerra JE, Hogue CJ, Atrash HK, Perez N. Infant mortality among Hispanics. A portrait of heterogeneity. *JAMA.* 1991;265:217–221.
108. Cohen BB, Friedman DJ, Mahan CM, Lederman R, Munoz D. Ethnicity, maternal risk, and birth weight among Hispanics in Massachusetts, 1987–89. *Public Health Rep.* 1993;108: 363–371.
109. Darabi KF, Ortiz V. Childbearing among young Latino women in the United States. *Am J Public Health.* 1987;77:25–28.
110. Erickson PI. Lessons from a repeat pregnancy prevention program for Hispanic teenage mothers in east Los Angeles. *Fam Plann Perspect.* 1994;26:174–178.
111. Kleinman JC. Infant mortality among racial/ethnic minority groups, 1983–1984. *MMWR CDC Surveill Summ.* 1990;39(SS-3):31–39.
112. Loria CM, Crespo CJ, Burt V. Blood pressure among Mexican-American, Cuban-American, and mainland Puerto Rican children. *Public Health Rep.* 1996;111(suppl 2):22–24.
113. Mendoza FS, Ventura SJ, Valdez RB, et al. Selected measures of health status for Mexican-American, mainland Puerto Rican, and Cuban-American children. *JAMA.* 1991;265:227–232.
114. Newacheck PW, Stoddard JJ, McManus M. Ethnolcultural variations in the prevalence and impact of childhood chronic conditions. *Pediatrics.* 1993;91:1031–1039.
115. Fact sheet. Healthy People 2000: Progress Review, Hispanic Americans. Washington, DC: US Department of Health and Human Services; 1997.
116. Wood PR, Hidalgo HA, Prihoda TJ, Kromer ME. Hispanic children with asthma: morbidity. *Pediatrics.* 1993;91:62–69.
117. Chabra A, Chávez GF, Harris ES. Mental illness in elementary-school-aged children. *West J Med.* 1999;170:28–34.
118. *Assuring Cultural Competence in Health Care: Recommendations for National Standards and an Outcomes-Focused Research Agenda.* Rockville, Md: US Department of Health and Human Services; 1999.
119. Hahn R, Mulinare J, Teutsch SM. Inconsistencies in coding of race and ethnicity between birth and death in US infants: a new look at infant mortality, 1983 through 1985. *JAMA.* 1992;267: 259–263.
120. *One Voice, One Vision—Recommendations to the Surgeon General to Improve Hispanic/Latino Health.* Washington, DC: US Surgeon General’s Hispanic/Latino Health Initiative; 1993.
121. *Hispanic Agenda for Action: Improving Services to Hispanic Americans.* Washington, DC: US Department of Health and Human Services Working Group on Hispanic Issues; 1996.
122. Guendelman S, Gould JB, Hudes M, Eskenazi B. Generational differences in perinatal health among the Mexican American population: findings from HHANES 1982–84. *Am J Public Health.* 1990;80(suppl):61–65.
123. Ismail A, Szpunar SM. The prevalence of total tooth loss, dental caries, and periodontal disease among Mexican Americans, Cuban Americans, and Puerto Ricans: findings from HHANES 1982–1984. *Am J Public Health.* 1990;80:66–70.
124. Lee DJ, Gomez-Marín O, Lee HM. Sociodemographic and educational correlates of hearing loss in Hispanic children. *Pediatr Perinat Epidemiol.* 1997;11:333–344.
125. Schwartz J, Otto D. Lead and minor hearing impairment. *Arch Environ Health.* 1991;46: 300–305.
126. Scribner R, Dwyer JH. Acculturation and low birthweight among Latinos in the Hispanic HHANES. *Am J Public Health.* 1989;79: 1263–1267.
127. Centers for Disease Control and Prevention. Update: blood lead levels—United States, 1991–1994. *MMWR Morb Mortal Wkly Rep.* 1997;46:141–146.
128. Pirkle JL, Brody DJ, Gunter EW, et al. The decline in blood lead levels in the United States: the National Health and Nutrition Examination Surveys (NHANES). *JAMA.* 1994;272: 284–291.
129. Guendelman S, English P, Chavez G. Infants of Mexican immigrants: health status of emerging populations. *Med Care.* 1995;33:41–52.
130. Massey DS, Zambrana RE, Alonzo-Bell S. Contemporary issues in Latino families: future directions for research, policy, and practice. In: Zambrana RE, ed. *Understanding Latino Families: Scholarship, Policy, and Practice.* Thousand Oaks, Calif: Sage Publications; 1995: 190–204.
131. Hayes-Bautista DE, Hurtado A, Valdez RB, Hernandez ACR. *No Longer a Minority: Latinos and Social Policy in California.* Los Angeles, Calif: Chicano Studies Research Center, University of California; 1992.
132. Szilagyi PG, Holl JL, Rodewald LE, et al. Evaluation of children’s health insurance: from New York State’s Child Health Plus to SCHIP. *Pediatrics.* 2000;105:687–691.
133. Health Care Financing Administration. Texas Title XXI fact sheet. Available at: <http://www.hcfa.gov/init/chpftx.htm>. Accessed August 14, 2000.
134. Immigration and Naturalization Service. Fact sheet: public charge. Available at: http://www.ins.usdoj.gov/graphics/publicaffairs/factsheets/public_cfs.htm. Accessed October 25, 2000.
135. Bachrach KM, Sandler IN. A retrospective assessment of inhalant abuse in the barrio: implications for prevention. *Int J Addict.* 1985;20: 1177–1189.
136. Smith JC, Mercy JA, Rosenberg ML. Suicide and homicide among Hispanics in the Southwest. *Public Health Rep.* 1986;101:265–270.
137. Holahan J, Wiener J, Wallin S. *Health Policy for the Low-Income Population: Major Findings From the Assessing the New Federalism Case Studies.* Washington, DC: Urban Institute; 1998.
138. Mexican American Study Project. *Health Status and Practices of Mexican Americans.* Los Angeles, Calif: Chicano Studies Research Center, University of California; 1968.
139. Furino A. *Health Policy and the Hispanic.* Boulder, Colo: Westview Press; 1992.